



ELSEVIER

Topology and its Applications 105 (2000) 113–119

TOPOLOGY
AND ITS
APPLICATIONS

www.elsevier.com/locate/topol

Topological and Borel compactifications of Polish G -spaces

Tzvi Scarr^{1,2}

Bar-Ilan University, Department of Mathematics and Computer Science, 52900 Ramat Gan, Israel

Received 11 December 1998; accepted 3 March 1999

Abstract

We investigate the relationships between topological and Borel G -spaces, where G is a Polish group. We show that every Polish G -space can be topologically and equivariantly embedded into a compact Polish G -space iff G is locally compact. This answers a question of Kechris. It also provides a striking contrast to the recent result of Becker and Kechris which states that every Borel G -space can be Borel-embedded into a compact Polish G -space. © 2000 Elsevier Science B.V. All rights reserved.

Keywords: Borel G -space; G -compactification; Game theory; Strong Choquet game

AMS classification: 54H15; 54H05; 22A99

1. Introduction

An *action* of a group G (with neutral element e) on a set X is a function $\alpha : G \times X \rightarrow X$ such that $\alpha(e, x) = x$ for all $x \in X$ and $\alpha(g_1 g_2, x) = \alpha(g_1, \alpha(g_2, x))$ for all $g_1, g_2 \in G$ and $x \in X$.

In topological dynamics, for example, G is a topological group, X is a topological space, and α is *continuous*. The triple $\langle G, X, \alpha \rangle$ is called a G -space, or a *topological G -space*. For basic information on topological G -spaces, see [20,25].

In descriptive set theory, one usually requires G to be a Polish (= complete, metrizable and separable) group and X a *standard Borel space*, that is, a measurable space (a set X and a σ -algebra \mathcal{S} of subsets of X) such that there exists a Polish topology on X with \mathcal{S} its σ -algebra of Borel sets. The assumption on the action α is weakened from continuity to Borel measurability. If X and Y are separable metrizable spaces, then a function $f : X \rightarrow Y$ is *Borel measurable* if $f^{-1}(B)$ is a Borel set in X for every Borel set B in Y . Thus, a *Borel*

¹ E-mail: scarr@avoda.jct.ac.il.

² This work is part of the author's Ph.D. Dissertation at Bar-Ilan University under the supervision of Prof. H. Furstenberg.

G -space is a triple $\langle G, X, \alpha \rangle$, where G is a Polish group, X is a standard Borel space, and $\alpha: G \times X \rightarrow X$ is a Borel-measurable action. See [3] for more on Borel G -spaces.

It is natural to compare topological and Borel G -spaces. What (kinds of) theorems are true in both contexts? How much flexibility is gained by relaxing the continuity of the action to Borel-measurability?

We have found a striking contrast in the area of G -space compactifications. A G -compactification of a topological G -space $\langle G, X, \alpha \rangle$ consists of a topological G -space $\langle G, cX, \alpha' \rangle$, where $\varphi: X \rightarrow cX$ is a compactification of X and $\alpha'|_{G \times X} = \alpha$. In other words, X is topologically and *equivariantly* embedded in cX , that is, $\varphi(gx) = g\varphi(x)$ for all $g \in G$ and $x \in X$.

A G -space is called G -Tychonoff if it has a G -compactification. For example, it is well known (see, for example, [4] and [17]) that for every topological group G , the G -space $\langle G, G, \alpha_L \rangle$ ($\alpha_L(g_1, g_2) = g_1 g_2$) is G -Tychonoff. In fact, $\langle G, G, \alpha_L \rangle$ has a maximal G -compactification, denoted by $\beta_G G$ and called the *greatest ambit*. Jan de Vries [21] proved that every coset G -space $\langle G, G/H, \alpha_L^* \rangle$ is G -Tychonoff, where $\alpha_L^*(g_1, g_2 H) = (g_1 g_2)H$. This includes the previous result as the particular case $H = \{e\}$. Ludescher and de Vries [10] showed that any G -space under an *equicontinuous* action is G -Tychonoff.

G is a V -group if every Tychonoff G -space is G -Tychonoff. In [21], Jan de Vries posed the “compactification problem” in its full generality: is every topological group G a V -group? Carlson [5] had earlier given a positive answer for the case $G = \mathbb{R}$. Palais [16] showed that every compact Lie group is a V -group. De Vries [22] and Antonyan [1] independently proved that every compact group is a V -group. In [24], de Vries improved these results.

Theorem 1.1 (de Vries). *Every locally compact group is a V -group.*

The converse of Theorem 1.1 is still open, and, in fact, it is not known whether the additive group \mathbb{Q} of rational numbers is a V -group. However, in [15] Megrelishvili and the present author showed that the class of non- V -groups is, in fact, rather large and contains all groups which are \aleph_0 -bounded (for example, second countable) and not locally precompact. Recall that a topological group G is called \aleph_0 -bounded [2,6] if for every neighborhood V of e there exists a countable subset S of G such that $SV = G$. Guran [6] proved that G is \aleph_0 -bounded iff G is a topological subgroup of a product of second countable topological groups. If G is separable, Lindelöf, or satisfies the countable chain condition, then G is \aleph_0 -bounded.

G is *locally precompact* if it is a subgroup of a locally compact group, or, equivalently, if its sup-completion (the completion with respect to its two-sided uniformity) is locally compact.

Theorem 1.2 [15]. *If G is an \aleph_0 -bounded topological group which is not locally precompact, then G is not a V -group.*

If G is Polish, we have a complete answer to the compactification problem.

Theorem 1.3 [15]. *Let G be a Polish group. Then G is a V -group iff G is locally compact.*

There is also much recent work in the area of *universal G -spaces*. In [7], Hjorth proves the existence of a *continuously universal* Polish G -space for every Polish group G . Recall also that for every second countable G and second countable G -Tychonoff X , there exists an equivariant topological embedding of the pair (G, X) into the pair $(H(I^{\aleph_0}), I^{\aleph_0})$, where I^{\aleph_0} denotes the Hilbert cube and $H(I^{\aleph_0})$ denotes the topological group of all homeomorphisms on the Hilbert cube. This follows as a particular case of Theorem 2.7 from [14].

Turning to Borel G -spaces, we find the notion of *universal Borel G -spaces*. Fix a Polish group G . A *universal Borel G -space* is a Borel G -space \mathcal{U}_G such that every Borel G -space can be Borel-embedded into \mathcal{U}_G . Mackey [11] and Varadarajan [19] proved that if G is Polish and *locally compact*, then there exists a universal Borel G -space. Becker and Kechris [3] have recently extended this result to *all* Polish groups, even ones which are *not* locally compact. In fact, they have gone even one step further.

Theorem 1.4 [3, Theorem 2.6.6]. *For any Polish group G , there is a universal Borel G -space which is moreover a compact Polish G -space.*

This last result means that for *every* Polish group G , every Borel G -space can be Borel-embedded into a compact Polish G -space.

Our Main Theorem, which was conjectured by Kechris [9], shows that the situation is strikingly different for *topological G -spaces*.

Main Theorem. *Let G be a Polish group. Then every Polish G -space can be (topologically) embedded into a compact Polish G -space iff G is locally compact.*

Our proof of the above theorem uses the *strong Choquet game* from descriptive set theory. We also need the following result.

Theorem 1.5 [13]. *Every G -Tychonoff space X has a G -compactification Y such that $w(Y) \leq w(X) \cdot w(G)$ and $\dim Y \leq \dim \beta_G X$.*

2. Proof of the Main Theorem

We turn now to the proof of the Main Theorem. We need a little background from descriptive set theory. See, for example, [8].

Fact 2.1. *A locally compact Hausdorff space is Polish iff it is second countable.*

We will use the following game.

Definition 2.2. Given a nonempty topological space X , the *strong Choquet game* G_X^s between two players is defined as follows:

Player I plays $x_0, U_0 \quad x_1, U_1 \quad \dots$
 Player II plays $V_0 \quad V_1$

Players I and II take turns playing nonempty open subsets of X such that $U_0 \supseteq V_0 \supseteq U_1 \supseteq V_1 \supseteq \dots$. Additionally, player I is required to play a point $x_n \in U_n$ and then player II must play $V_n \subseteq U_n$ with $x_n \in V_n$.

Player I wins the game if $\bigcap_n V_n = \emptyset$.

Player II wins the game if $\bigcap_n V_n (= \bigcap_n U_n) \neq \emptyset$.

A nonempty space X is called a *strong Choquet space* if player II has a winning strategy for the game G_X^s .

Theorem 2.3 (Choquet). *A nonempty, second countable topological space is Polish iff it is T_1 , regular, and strong Choquet.*

We can now prove the Main Theorem.

Main Theorem 2.4. *Let G be a Polish group. Then every Polish G -space can be (topologically) embedded into a compact Polish G -space iff G is locally compact.*

Proof. (\Leftarrow) Let X be a Polish G -space. Since G is locally compact, X has a G -compactification, by Theorem 1.1. Since G and X are both second countable, there exists a second countable G -compactification Y of X , by Theorem 1.5. By Fact 2.1, Y is Polish.

(\Rightarrow) Suppose G is Polish and not locally compact. Then G is second countable and not locally precompact. We will construct a Polish G -space which is not G -Tychonoff. This construction can also be found in [15].

Since G is not locally precompact, G does not act locally uniformly equicontinuously on $X = \beta_G G$ (see [15, Lemma 2.7]). We will first construct a G -space X_U for every U in a collection \mathbb{B} (of cardinality $\chi(G)$) of basic neighborhoods of e in G . By [15, Fact 2.8], X is a G -limit of an inverse G -system of compact metrizable G -spaces X_i ($i \in I$). Let μ and μ_i denote the unique compatible uniformity on X and X_i , respectively.

Let $U \in \mathbb{B}$. Since U does not act μ -uniformly equicontinuously on $\beta_G G$, there exists an index $i \in I$ such that U does not act μ_i -uniformly equicontinuously on (X_i, μ_i) . Therefore, there is $\varepsilon \in \mu_i$ such that for every $\delta \in \mu_i$ there exist $(x_\delta, y_\delta) \in \delta$ and $g_\delta \in U$ such that

$$(g_\delta x_\delta, g_\delta y_\delta) \notin \varepsilon. \quad (*)$$

Thus we obtain nets $\langle x_\delta \rangle$, $\langle y_\delta \rangle$, $\langle g_\delta x_\delta \rangle$, and $\langle g_\delta y_\delta \rangle$ in (X_i, μ_i) , indexed by the elements δ of μ_i . Passing to subnets if necessary, we may assume that there exist $x^U, a^U, b^U \in X$ such that $x_\delta \rightarrow x^U$, $y_\delta \rightarrow x^U$, $g_\delta x_\delta \rightarrow a^U$, and $g_\delta y_\delta \rightarrow b^U$. By (*) we have $a^U \neq b^U$. Hence $(x^U, x^U) \in \Delta$, and $(a^U, b^U) \notin \Delta$, where $\Delta = \{(x, x) \mid x \in X_i\}$.

We put the “two-coordinate” action on the G -space $(X_i, \mu_i) \times (X_i, \mu_i)$ and then form the quotient G -space $Y_i = (X_i \times X_i)/\Delta$. Consider the quotient G -map $p: X_i \times X_i \rightarrow Y_i$. Let $z := p(x^U, x^U)$.

Next, let

$$X_U = (Y_i \times Y_i) \setminus \{(z, z)\}, \quad C^U = (\{z\} \times Y_i) \setminus \{(z, z)\},$$

and

$$D^U = (Y_i \times \{z\}) \setminus \{(z, z)\}.$$

Define the “one-coordinate” action $\alpha': G \times X_U \rightarrow X_U$ by

$$\alpha'(g, (x, y)) = (\alpha(g, x), y),$$

where α is the action of G on Y_i . This completes the construction of X_U from U .

Now form the topological G -sum $S = \bigoplus \{X_U: U \in \mathbb{B}\}$. Let $\alpha^*: G \times S \rightarrow S$ be the natural action. Define

$$C = \bigcup_{U \in \mathbb{B}} C^U, \quad D = \bigcup_{U \in \mathbb{B}} D^U.$$

Finally, from $\langle G, S, \alpha^* \rangle$, we will construct a normal G -space $\langle G, S^+, (\alpha^*)^+ \rangle$ which is not G -Tychonoff.

Let $\omega = \mathbb{N} \cup \{0\}$ carry the discrete topology. Let Y be the quotient space formed from $S \times \omega$ by identifying the pairs $(c, 2i + 1)$ and $(c, 2i + 2)$ for $c \in C, i \in \omega$, and the pairs $(d, 2i)$ and $(d, 2i + 1)$ for $d \in D, i \in \omega$. Let $p: S \times \omega \rightarrow Y$ be the quotient map. For $n \in \omega$, let $i_n: S \rightarrow S \times \omega$ be the canonical injection $x \mapsto (x, n)$.

Fix a point $a \notin Y$, and let $S^+ = Y \cup \{a\}$. Topologize S^+ by setting Y to be an open subset with its quotient topology and the n th basic neighborhood of a to be

$$N_n(a) = \{a\} \cup p(i_{2n}(S \setminus C)) \cup \bigcup \{p(i_m(S)): m > 2n\}.$$

We now define $\alpha^+: G \times S^+ \rightarrow S^+$. For any $g \in G$, set $\alpha^+(g, a) = a$. For $p((x, n)) \in Y$, set $\alpha^+(g, p((x, n))) = p(i_n(\alpha^*(g, x)))$. Then $\langle G, S^+, \alpha^+ \rangle$ is a normal G -space which is not G -Tychonoff (see [15, Theorem 3.2]).

From the construction of S and S^+ , it is clear that if G is second countable, then S^+ can be chosen to be second countable. We will show that S^+ is, in fact, a Polish space. Since S^+ is normal, it is enough, by Theorem 2.3, to show that S^+ is strong Choquet.

Let $Y = S^+ \setminus \{a\}$. Then Y is locally compact, Hausdorff, and second countable. By Fact 2.1, Y is Polish. So Y is strong Choquet, by Theorem 2.3. Now we can define a winning strategy for player II in the game $G_{S^+}^s$.

Suppose player I plays x_0, U_0 . If $a \notin U_0$, then U_0 is an open subset of Y . So player II can now apply his winning strategy for the game G_Y^s .

Otherwise, $a \in U_0$. If $x_0 \neq a$, then player II can play $U_0 \cap Y$. This will force player I to play x_1, U_1 , where U_1 is an open subset of Y . So player II can now apply his winning strategy for G_Y^s .

Finally, suppose $x_0 = a$. Player II can play $V_0 = N_n(a)$, where $n \in \mathbb{N}$ is large enough so that $N_n(a) \subseteq U_0$. From this point on, if player I plays x_k, U_k , with $U_k \subseteq Y$, then player II

applies his winning strategy for G_Y^s . Otherwise, player I always plays a , U_k . In this case, player II can play $V_k = N_{n_k}(a)$ for sufficiently large and increasing values of n_k . In the end, we will have

$$\bigcap_k V_k = \{a\},$$

and player II wins the game $G_{S^+}^s$.

Therefore S^+ is strong Choquet and, hence, Polish. \square

3. Question

When can a Tychonoff G -space be topologically embedded into a compact *Borel* G -space?

Acknowledgements

The author is very grateful to his advisor, Professor Hillel Furstenberg. He would also like to thank Professor Alexander Kechris for the question and for stimulating conversations and Professor Gerald Itzkowitz for his helpful comments.

References

- [1] S. Antonyan, Yu. Smirnov, Universal objects and bicomact extensions for topological groups of transformations, *Soviet Math. Dokl.* 23 (1981) 279–284.
- [2] A.V. Arhangel'skiĭ, Classes of topological groups, *Russian Math. Surveys* 36 (1981) 521–526.
- [3] H. Becker, A. Kechris, *The Descriptive Set Theory of Polish Group Actions*, Cambridge Univ. Press, Cambridge, 1996.
- [4] R.B. Brook, A construction of the greatest ambit, *Math. Systems Theory* 4 (1970) 243–248.
- [5] D.H. Carlson, Extensions of dynamical systems via prolongations, *Funkcial. Ekvac.* 14 (1971) 35–46.
- [6] I.I. Guran, On topological groups close to being Lindelöf, *Soviet Math. Dokl.* 23 (1981) 173–175.
- [7] G. Hjorth, A universal Polish G -space, *Topology Appl.* 91 (1999) 141–150.
- [8] A. Kechris, *Classical Descriptive Set Theory*, Springer, New York, 1995.
- [9] A. Kechris, e-mail communication, October, 1997.
- [10] H. Ludescher, J. de Vries, A sufficient condition for the existence of a G -compactification, *Nederl. Akad. Wetensch. Proc. Ser. A* 83 (1980) 263–268.
- [11] G.W. Mackey, Point realizations of transformation groups, *Illinois J. Math.* 6 (1962) 327–335.
- [12] M.G. Megrelishvili, Equivariant completions and compact extensions, *Bull. Acad. Sci. Georgian SSR* 115 (1984) 21–24.
- [13] M. Megrelishvili, Compactification and factorization in the category of G -spaces, in: J. Adámek, S. MacLane (Eds.), *Categorical Topology and its Relation to Analysis, Algebra and Combinatorics*, World Scientific, Singapore, 1989, pp. 220–237.
- [14] M. Megrelishvili, Free topological G -groups, *New Zealand J. Math.* 25 (1996) 59–72.
- [15] M. Megrelishvili, T. Scarr, Constructing Tychonoff G -spaces which are not G -Tychonoff, *Topology Appl.* 86 (1998) 69–81.

- [16] R. Palais, The classification of G -spaces, *Mem. Amer. Math. Soc.* 36 (1960) 25.
- [17] S. Teleman, Sur la représentation linéaire des groupes topologiques, *Ann. Sci. Ecole Norm. Sup.* 74 (1957) 319–339.
- [18] I. Tree, Constructing regular spaces that are not completely regular, *Houston J. Math.* 21 (3) (1995) 613–622.
- [19] V.S. Varadarajan, Groups of automorphisms of Borel spaces, *Trans. Amer. Math. Soc.* 109 (1963) 191–220.
- [20] J. de Vries, *Topological Transformation Groups I*, Math. Centre Tracts 65, Mathematisch Centrum, Amsterdam, 1975.
- [21] J. de Vries, Can every Tychonoff G -space equivariantly be embedded in a compact Hausdorff G -space?, *Math. Centrum, Amsterdam, Afd. Zuivere Wisk.* 36 (1975).
- [22] J. de Vries, A note on compactifications of G -spaces, *Math. Centrum, Amsterdam, Afd. Zuivere Wisk.* 61 (1976).
- [23] J. de Vries, Equivariant embeddings of G -spaces, in: *General Topology and its Relations to Modern Analysis and Algebra IV, Part B*, Proc. 4th Prague Topological Symposium (1976), Prague (1977) 485–493.
- [24] J. de Vries, On the existence of G -compactifications, *Bull. Acad. Polon. Sci. Ser. Math.* 26 (1978) 275–280.
- [25] J. de Vries, *Elements of Topological Dynamics*, Kluwer Academic Publishers, Dordrecht, 1993.